



## Corrigendum

**Corrigendum to “Photocatalytic carbon oxidation with nitric oxide”  
[Appl. Catal. B: Environ. 166–167 (2015) 374–380]**

Lijun Liao<sup>a</sup>, Steven Heylen<sup>a</sup>, Brecht Vallaey<sup>a</sup>, Maarten Keulemans<sup>a,b</sup>, Silvia Lenaerts<sup>b</sup>,  
Maarten B.J. Roefsaers<sup>a</sup>, Johan A. Martens<sup>a,\*</sup>

<sup>a</sup> Centre for Surface Chemistry and Catalysis, KU Leuven, Kasteelpark Arenberg 23, 3001 Heverlee, Belgium

<sup>b</sup> Research Group Sustainable Energy and Air Purification, Department of Bioscience Engineering, University of Antwerp, Groenenborgerlaan 171, B-2020 Antwerp, Belgium

The authors regret to inform that the numerical values of the carbon oxidation rates were incorrect.

In the long experiment of Fig. 4 the carbon oxidation rate was ca. 0.4  $\mu\text{g}$  carbon per hour and per mg  $\text{TiO}_2$ . The highest carbon oxidation rate of ca. 2  $\mu\text{g}$  carbon per hour and per mg  $\text{TiO}_2$  was obtained in the experiment with 3000 ppm NO in dry gas, corresponding to ca. 2.3  $\text{nmol s}^{-1}$  (Fig. 2). Photooxidation of carbon particles emitted by a light duty vehicle would require about 2.9 kg of  $\text{TiO}_2$ .

Authors would like to apologize for the inconvenience caused.

DOI of original article: <http://dx.doi.org/10.1016/j.apcatb.2014.11.051>.

\* Corresponding author. Tel.: +32 16 32 16 37; fax: +32 16 32 19 98.

E-mail address: [johan.martens@biw.kuleuven.be](mailto:johan.martens@biw.kuleuven.be) (J.A. Martens).